Ecology of Diamondback Terrapins in North Inlet, Winyah Bay, SC

Peter King, FMU
Jennifer Spicer, NERR
Paige Weaver, FMU










Aims


- Mark recapture population study
- Size distribution -population viability
- Habitat utilization feeding, mating, nesting, juveniles
- Site fidelity
- Diet












Data

- 9 creeks surveyed in summer 2006
- Terrapins found in 5 out of 9
- 45 terrapins, 20 female, 25 male
- 11 creeks surveyed in summer 2007
- Terrapins found in 9 out 11
- 45 terrapins, 15 female, 30 male
- Total
 86 terrapins, 33 female, 53 male

Recaptures

- 4 recaptures
- 3 from same creek
- 1 from neighboring creek





90 terrapin captures in 14 creeks

2 recaptures from OM, E5

2 recaptures, 1 from OM, E4 1 from OM, E5

Frequency of males and females by size class






Confirmed nesting areas



Evidence of nesting activity

May 30 - June 13





















Conclusions to date

- Distribution of sizes indicates recruitment
- Little known about juveniles
- High incidence of nest depredation
- No evidence of crab trap mortality
- Recommend nest site protection and predator control

Thanks to the following for help

Karen Philips

Pat Cloninger

Lacy Danikas

Travis Knowles

Jeff Camper

Kirk Dineley

Nicole Reuter

Philip Vernon

Tony Arthur

Edward Arnold

Rebecca Koontz

Jeremy Capps

Anton Capers

Stephanie Miller

Whit Gibbons et al

Dennis Allen et al

(BFML)



Funding and assistance FMU, Baruch Foundation, BMFL